

EFFECTS OF PROLONGED EXPOSURE TO CIGARETTE SMOKE ON PHENOTYPE AND FUNCTION OF INNATE IMMUNE CELLS IN THE LUNGS

Vladislav Volarević¹, Nikolina Kastratović¹, Ana Volarević²

Cigarette smoke contains various toxic compounds that can alter the phenotype and function of immune cells within the lungs. To investigate how extended exposure to cigarette smoke affects innate immune cells (including neutrophils, macrophages, dendritic cells (DCs), natural killer (NK) cells, and natural killer T (NKT) cells), we exposed BALB/c mice to cigarette smoke for 4 or 8 weeks, with control animals being exposed to air. The cigarette smoke was generated using an LM1 Borgwaldt smoking machine. Flow cytometry analysis and intracellular staining of lung-infiltrated immune cells revealed that prolonged exposure to cigarette smoke notably increased the number of inflammatory cluster of differentiation (CD14) receptor-expressing neutrophils, tumor necrosis factor- α (TNF- α)-producing NK and NKT cells, and inflammatory cluster of differentiation (CD40)-expressing macrophages. Additionally, significantly higher numbers of interleukin-12 (IL-12) and interleukin-23 (IL-23) producing DCs were observed in the lungs of mice exposed to cigarette smoke for 8 weeks compared to those exposed to air. Moreover, a greater number of TNF- α , IL-12, and IL-23-producing macrophages, along with TNF- α -producing neutrophils, NK, and NKT cells, were found in the lungs of mice exposed to cigarette smoke for 8 weeks compared to those exposed for 4 weeks. In conclusion, prolonged cigarette smoke exposure promotes an inflammatory phenotype in lung-infiltrating innate immune cells, contributing to the progression and worsening of inflammatory lung diseases.

Acta Medica Medianae 2025;64(4):145–152.

Key words: cigarette smoke, lungs, macrophages, neutrophils, dendritic cells

¹University of Kragujevac, Faculty of Medical Sciences, Department of Genetics, Microbiology and Immunology, Center for Research on Harmful Effects of Biological and Chemical Hazards, Kragujevac, Serbia

²University of Kragujevac, Faculty of Medical Sciences, Department of Psychology, Center for Research on Harmful Effects of Biological and Chemical Hazards, Kragujevac, Serbia

Contact: Vladislav Volarević
E-mail: vladislav.volarevic@medfak.ni.ac.rs
69 Svetozara Markovica St., 34000 Kragujevac, Serbia

Introduction

Cigarette smoke is composed of numerous harmful substances that severely impact the respiratory system and play a major role in the onset and advancement of lung-related diseases (1–5). Formaldehyde, acrolein, benzene, and heavy metals cause oxidative stress-induced damage of alveolar epithelial cells, resulting in structural changes in the lungs (1, 2). Continuous and long-term exposure to these harmful components can lead to the destruction of alveolar walls, resulting in the development of emphysema (3, 4). Additionally, alarmins released from injured alveolar epithelial cells can recruit fibroblasts and

trigger excessive collagen deposition, leading to the development of pulmonary fibrosis (4). Moreover, as highly carcinogenic compounds, polycyclic aromatic hydrocarbons from cigarette smoke can cause mutations in lung parenchymal cells, paving the way for the development of lung cancer (5).

Cigarette smoke significantly affects the microbiome of the respiratory tract, leading to the development of dysbiosis, an imbalance in the microbial communities within the lungs (6, 7). Long-term exposure to the harmful chemicals of cigarette smoke creates a hostile environment for beneficial bacteria while promoting the growth of pathogenic microorganisms (6, 7). This dysbiotic state can impair the lungs' ability to defend against microbial pathogens and may induce a detrimental immune response in the lungs (6, 7). The inflammatory response triggered by the altered microbiome could create a vicious cycle, in which the inflammation promotes the persistence of harmful microbes, exacerbating lung damage and contributing to the development and progression of chronic pulmonary diseases (8).

Importantly, harmful components of cigarette smoke may disrupt the normal homeostasis of the immune system in the lungs, enabling the development of immune cell-driven inflammatory diseases (9). An altered expression

of adhesion molecules (selectins and integrins) was observed on the endothelial cells that were continuously exposed to cigarette smoke (10). Accordingly, prolonged exposure to cigarette smoke affects the migration of circulating immune cells in the lungs, impairing their capacity to elicit an appropriate anti-microbial immune response (9). Additionally, harmful components of cigarette smoke could trigger a potent, detrimental immune response in the lungs by inducing the massive release of pro-inflammatory cytokines from lung tissue-resident innate immune cells (natural killer (NK), natural killer T (NKT) cells, macrophages, and dendritic cells (DCs)), setting off a cycle of chronic inflammation (11). Continuously activated innate immune cells orchestrate detrimental inflammatory response by promoting activation of adaptive immune cells, T and B lymphocytes (11, 12). Persistent activation of immune cells within the lungs over an extended period leads to significant tissue injury and structural changes in the airways, ultimately contributing to the emergence of chronic inflammatory lung conditions such as asthma and chronic obstructive pulmonary disease (COPD) (9, 13).

In order to better understand how prolonged exposure to cigarette smoke affects lung neutrophils, macrophages, DCs, NK, and NKT cells, we herewith used a well-established murine model of lung inflammation induced by cigarette smoke (3, 11) to examine the variations in phenotype and cytokine production of these innate immune cells following one and two months of cigarette smoke exposure.

Materials and Methods

Experimental Animals

This study involved BALB/c mice aged six to eight weeks. The animals were kept under specific-pathogen-free conditions in a climate-regulated facility, featuring a 12-hour alternating light/dark cycle, at the Animal Facility of the Faculty of Medical Sciences, University of Kragujevac, Serbia. They were provided unlimited access to clean water and a standard diet formulated for laboratory rodents. All procedures involving the animals complied with ethical standards and were approved by the Animal Ethics Committee of the same institution, ensuring the welfare of the animals throughout the study.

Design of the Study

The mice were randomly assigned to three groups: a control group exposed only to ambient air, and two experimental groups subjected to cigarette smoke for either 4 weeks (Group 1) or 8 weeks (Group 2). Cigarette smoke was produced using Marlboro Red (MR) cigarettes and delivered via an LM1 Borgwaldt smoking apparatus. Exposure followed the Health Canada Intense (HC1) puffing protocol, which involves drawing

nine puffs per cigarette with each puff lasting 2 seconds and delivering 55 mL of smoke in a bell-shaped flow profile (27.5 mL/s), taken every 30 seconds with blocked filter vents, down to a butt length of 36 ± 0.5 mm. Before use, MR cigarettes were conditioned for a minimum of 48 hours at $60 \pm 3\%$ relative humidity and $22 \pm 1^\circ\text{C}$. Smoke was administered in an environment maintained at $60 \pm 5\%$ relative humidity and $22 \pm 2^\circ\text{C}$, as per ISO 3402:1999 standards (11). The animals were euthanized after completing the exposure protocol, specifically 24 hours following their final exposure to cigarette smoke.

Isolation of Immune Cells From the Mice Lungs

Initially, the mice lungs underwent enzymatic digestion. Once removed, the lungs were placed in a petri dish, rinsed with phosphate-buffered saline (PBS), and finely minced using a scalpel or razor blade. The resulting tissue fragments were transferred into 15 mL conical tubes containing a digestion mixture composed of Dulbecco's Modified Eagle Medium (DMEM) supplemented with 0.5 mg/mL collagenase IV and 1 mg/mL DNase. The samples were incubated at 37°C in a water bath for 45 minutes with continuous agitation at 80 rpm. After digestion, the cell suspensions were first passed through 100 μm nylon mesh filters into 50 mL tubes containing DMEM enriched with 10% fetal bovine serum (FBS), followed by centrifugation. The resulting cell pellets were then separated from the supernatant and treated with 5 mL of lysis buffer for 5 minutes to remove red blood cells. A subsequent centrifugation step was performed, after which the cells were resuspended in fresh medium and prepared for intracellular staining and flow cytometric evaluation (11).

Flow Cytometry Analysis and Intracellular Staining of Immune Cells

Immune cells extracted from the lungs of mice exposed to either ambient air or cigarette smoke were assessed for a range of surface and intracellular markers using flow cytometry (11). To reduce potential cell loss or damage resulting from tissue dissociation processes, viable cells were selectively isolated using the MACS[®] Dead Cell Removal Kit (Miltenyi Biotec, Bergisch Gladbach, Germany; Cat. No. 130-090-101). For this process, single-cell suspensions were incubated with 100 μL of Dead Cell Removal MicroBeads per 10 million cells at room temperature for 15 minutes. The mixture was then passed through MS columns pre-filled with MACS Binding Buffer, allowing live cells to pass through while dead cells were retained.

To prevent nonspecific interactions with Fc receptors, the isolated live cells were pre-treated with anti-mouse cluster of differentiation (CD) 16 and 32 (CD16/CD32) (BD Fc Block), using 1 μg per

million cells in 100 μ L of staining buffer composed of Dulbecco's PBS (DPBS) without calcium or magnesium, 1% heat-inactivated fetal bovine serum (FBS), and 0.09% sodium azide. The incubation was carried out for 15 minutes at 4°C. After blocking, the cells were washed and stained with fluorescently labeled antibodies.

A total of 1×10^6 cells were incubated for 30 minutes in the dark at 4°C with antibodies targeting CD45, CD3, CD4, CD8, F4/80, CD11c, Ly6G/C, CD49, CD14, CD40, and TLR-2. These antibodies were conjugated with various fluorophores, including FITC, PE, PerCP, and APC (BD Biosciences, San Jose, CA, USA). The staining process concluded with two washing steps using staining buffer, followed by centrifugation to pellet the cells for subsequent analysis.

For intracellular marker detection, cells were activated using 50 ng/mL phorbol 12-myristate 13-acetate (PMA) and 500 ng/mL ionomycin for a duration of 5 hours. After this stimulation period, GolgiStop (BD Biosciences, San Jose, CA, USA; Cat. No. 554715) was introduced to inhibit protein transport. Cells were subsequently fixed and permeabilized using the BD Cytotfix/Cytoperm™ Fixation/Permeabilization Kit (Cat. No. 554714), with incubation at 4°C for 20 minutes. Following fixation, cells were rinsed twice with BD Perm/Wash™ buffer and centrifuged.

The permeabilized cells were then resuspended in 50 μ L of BD Perm/Wash™ buffer containing pre-titrated fluorochrome-conjugated antibodies targeting inducible nitric oxide synthase (iNOS), and pyrin domain-containing protein 3 (NLRP3), tumor necrosis factor- α (TNF- α), interleukin-12 (IL-12), and interleukin-23 (IL-23). These antibodies, specific for mouse antigens, were labeled with FITC, PE, PerCP, or APC fluorophores (BD Biosciences, San Jose, CA, USA). Staining was carried out at 4°C in the dark for 30 minutes. Afterward, cells underwent two additional washes with Perm/Wash™ buffer and were finally resuspended in staining buffer for flow cytometric analysis. Data collection was conducted using a BD FACSCalibur system, and results analyzed using Flowing software (Turku Bioscience Centre).

Statistical Analysis

All statistical evaluations were performed using SPSS software version 21.0 for Windows (SPSS, Inc., Chicago, IL, USA). To determine variations in group means, a one-way analysis of variance (ANOVA) was employed. In cases where ANOVA revealed statistically significant differences, *post hoc* comparisons between specific groups were conducted using two-tailed Student's t-tests, applying the Tukey method to adjust for multiple comparisons. Results are expressed as mean values \pm standard error of the mean (SEM). A p-value less than 0.05 was regarded as indicating statistical significance, and all tests were two-tailed.

Results

Extended Cigarette Smoke Exposure Promotes Accumulation of Inflammatory Neutrophils, Natural Killer and Natural Killer T Cells in Lung Tissue

Flow cytometric evaluation of immune cells isolated from the lungs demonstrated that chronic exposure to cigarette smoke markedly elevated the levels of inflammatory neutrophils, natural killer (NK) cells, and natural killer T (NKT) cells in the lungs of treated mice (Figure 1). A substantial increase in CD14⁺CD45⁺Ly6G⁺ neutrophil populations was observed in the lungs of mice exposed to smoke for 4 weeks (Figure 1A; $p < 0.01$) and even more so after 8 weeks of exposure (Figure 1A; $p < 0.001$), in comparison to the air-exposed control group. In addition, cigarette smoke inhalation for 8 weeks significantly enhanced the presence of neutrophils expressing TLR-2 (Ly6G⁺CD45⁺) in the lung tissue (Figure 1B; $p < 0.01$).

Further intracellular staining revealed that ongoing smoke exposure drove these neutrophils toward a more pro-inflammatory profile (Figure 1C-D). Mice exposed to cigarette smoke for 8 weeks exhibited a markedly higher number of inducible nitric oxide synthase (iNOS)-positive neutrophils (Figure 1C; $p < 0.001$), as well as TNF- α -secreting CD45⁺Ly6G⁺ neutrophils (Figure 1D; $p < 0.01$), compared to those exposed for only 4 weeks.

Similarly, prolonged exposure to cigarette smoke resulted in an increased presence of inflammatory TNF- α -producing CD49b⁺ NK cells (Figure 1E) and CD49b⁺CD3⁺ NKT cells (Figure 1F) in the lungs of mice. Significantly higher numbers of TNF- α -producing NK and NKT cells were detected in the lung tissue of mice exposed to cigarette smoke for both 4 weeks (Figure 1E-F; $p < 0.01$) and 8 weeks (Figure 1E-F; $p < 0.05$), compared to the control group exposed only to air. Notably, extended cigarette smoke exposure notably increased TNF- α production by both NK and NKT cells in the lungs, with the 8-week exposure group showing a significantly higher proportion of TNF- α -producing CD49b⁺ and CD49b⁺CD3⁺ cells when compared to the 4-week exposure group (Figure 1E-F; $p < 0.05$).

Flow cytometric analysis of immune cells in the lungs revealed that BALB/c mice exposed to cigarette smoke for 8 weeks had the highest levels of CD45⁺Ly6G/C⁺ neutrophils expressing CD14 (A), TLR-2 (B), iNOS (C), and TNF- α (D), as well as the highest numbers of TNF- α -producing NK (E) and NKT cells (F), compared to mice exposed to cigarette smoke for 4 weeks or air-exposed control mice. Data are shown as Mean \pm SEM; $n = 6$ mice per group; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Figure 1. The impact of extended cigarette smoke exposure on the presence of neutrophils, NK, and NKT cells in the lungs of mice

Chronic Cigarette Smoke Exposure Led to the Development of an Inflammatory Profile in Lung Macrophages

Analysis of lung cell populations revealed that extended exposure to cigarette smoke significantly increased the number of inflammatory macrophages (Figure 2). The presence of NLRP3 inflammasome-activated F4/80+ macrophages was notably higher in mice exposed to smoke for 4 weeks (Figure 2A; $p < 0.05$) and 8 weeks (Figure 2A; $p < 0.001$), compared to the control group exposed to air. Additionally, after 8 weeks of smoke exposure, there was a marked increase in the number of F4/80+ macrophages expressing iNOS (Figure 2B; $p < 0.001$), CD40 (Figure 2C; $p < 0.05$), and CD14 (Figure 2D; $p < 0.05$) in the lungs of the experimental animals.

Intracellular staining further demonstrated that prolonged exposure to cigarette smoke induced the production of pro-inflammatory cytokines such as IL-12, IL-23, and TNF- α in lung macrophages (Figure 2D-F). Significantly elevated levels of IL-12 (Figure 2D; $p < 0.001$), IL-23 (Figure 2E; $p < 0.001$), and TNF- α -producing macrophages (Figure 2F; $p < 0.05$) were detected in the lungs of mice exposed to cigarette smoke for 4 weeks, compared to those exposed to air. This effect was more pronounced in animals exposed to smoke for 8 weeks, with substantially higher numbers of IL-12 (Figure 2D; $p < 0.001$), IL-23 (Figure 2E; $p < 0.001$), and TNF- α -producing F4/80+ macrophages (Figure 2F; $p < 0.05$) observed in the 8-week exposure group compared to the 4-week exposure group.




Figure 2. The lasting impact of cigarette smoke on the phenotype and functionality of lung macrophages

A notable increase in the total number of F4/80+ macrophages expressing NLRP3 inflammasome (A), iNOS (B), CD40 (C), CD14 (D), and producing IL-12 (E), IL-23 (F), and TNF- α (G) was observed in BALB/c mice exposed to cigarette smoke for 4 or 8 weeks, compared to the air-exposed control group. Data are presented as Mean \pm SEM; n = 6 mice per group; *p < 0.05, **p < 0.01, ***p < 0.001.

Chronic Cigarette Smoke Exposure Increased the Number of Inflammatory DCs in the Lungs of Mice

Similar to the effects seen in macrophages, sustained exposure to cigarette smoke promoted the development of an inflammatory phenotype in dendritic cells (DCs) in the lungs (Figure 3). A significant rise in the numbers of IL-12 and IL-23 producing CD11c+ DCs (Figure 3A-B; p < 0.05) was observed in the lungs of mice exposed to smoke for 8 weeks, compared to the air-exposed control group. Additionally, the total count of NLRP3+CD11c+ dendritic cells (DCs) was significantly higher in the lungs of mice exposed to cigarette smoke for both 4 weeks and 8 weeks (Figure 3C; p < 0.001), compared to the control group.

The substantial increase in inflammatory DCs in the lungs coincided with a significant rise in the number of inflammatory T cells infiltrating the

lungs (Figure 3D-E). Notably, the number of TNF- α -producing helper CD4+ T cells was significantly higher in mice exposed to cigarette smoke for 4 weeks (Figure 3D; p < 0.01) and 8 weeks (Figure 3D; p < 0.001). Similarly, a marked increase in TNF- α -producing cytotoxic CD8+ T cells was observed in the lungs of mice exposed to cigarette smoke for both 4 weeks (Figure 3E; p < 0.05) and 8 weeks (Figure 3E; p < 0.001).

Crucially, prolonged cigarette smoke exposure significantly enhanced TNF- α production in both helper and cytotoxic T lymphocytes, as evidenced by a higher number of TNF- α -producing CD4+ (Figure 3D; p < 0.05) and CD8+ T cells (Figure 3E; p < 0.01) in the lungs of mice exposed to smoke for 8 weeks, compared to those exposed for 4 weeks.

Flow cytometry analysis of immune cells infiltrating the lungs showed that the highest numbers of IL-12 (A) and IL-23 producing (B) NLRP3 inflammasome-activated (C) CD11c+DCs, as well as TNF- α -producing CD4+ T helper (D) and cytotoxic CD8+ T cells (E), were found in the lungs of BALB/c mice exposed to cigarette smoke for 8 weeks, compared to those exposed for 4 weeks or the air-exposed control group. Data are presented as Mean \pm SEM; n = 6 mice per group; *p < 0.05, **p < 0.01, ***p < 0.001.




Figure 3. The effect of extended cigarette smoke exposure on the presence of DCs in the lungs

Discussion

Cigarette smoke exposure leads to an increased influx of neutrophils into the lungs, where they become activated and release various pro-inflammatory cytokines and chemokines (14, 15). These smoke-activated neutrophils generate reactive oxygen species (ROS) and proteolytic enzymes, which can damage lung tissue and worsen chronic inflammatory conditions such as COPD and asthma (14). Moreover, cigarette smoke alters neutrophil functions by suppressing apoptosis, thereby prolonging their survival in the lungs and contributing to ongoing inflammation (15). In our study, we observed that prolonged exposure to cigarette smoke enhanced the inflammatory response of neutrophils in the lungs. Specifically, long-term smoke exposure significantly increased the number of CD14 and TLR-2-expressing neutrophils, which play a crucial role in initiating and exacerbating lung inflammation. These receptors facilitate the detection of pathogen-associated molecular patterns, leading to neutrophil activation and a surge in ROS and inflammatory cytokine production, which further damages lung tissue (16). Consistent with these findings, we noted a significant increase in the number of iNOS-expressing and TNF- α -producing neutrophils in the lungs of mice exposed to cigarette smoke for 8 weeks.

Furthermore, our data showed that prolonged exposure to cigarette smoke led to an elevated presence of TNF- α -producing NK and NKT cells in the lungs. NK cells are essential for recognizing and eliminating infected or transformed cells through cytotoxic mechanisms (17), while NKT cells, which share features of both T and NK cells, contribute significantly to the development and progression of lung inflammation (17). Upon activation, these cells release large amounts of TNF- α , which induces the expression of adhesion molecules such as E and P selectins on endothelial cells, thereby promoting the recruitment of circulating leukocytes into the lungs (8, 17). The increase in TNF- α -producing NK and NKT cells due to cigarette smoke exposure may play a critical role in driving tissue damage and worsening inflammatory lung diseases. Chronic inflammation caused by these cells can lead to the destruction of alveolar structures, which is a key factor in the development of emphysema in lungs exposed to cigarette smoke over time (9).

Macrophages play a crucial role in the immune defense of the lungs, participating in phagocytosing pathogens, antigens presentation, and the modulation of inflammatory responses (18). When exposed to cigarette smoke, a variety of toxic chemicals are introduced that can alter their function and characteristics, leading to an imbalance in the macrophage-driven immune response (19). Macrophages exposed to cigarette smoke show reduced phagocytic activity, weakening their ability to eliminate inhaled

microbial threats (19). This impairment is partly due to the oxidative stress caused by cigarette smoke, which interferes with the signaling pathways required for macrophage activation and phagocytosis (19, 20). In our study, we found that prolonged exposure to cigarette smoke led to a higher number of macrophages with an activated NLRP3 inflammasome and increased expression of iNOS.

Additionally, cigarette smoke affects the ability of lung macrophages to present antigens effectively (11, 19). Exposure to smoke can alter the expression of co-stimulatory molecules, which hampers their capacity to present antigens to T cells, potentially leading to ineffective T cell activation and contributing to T cell-mediated lung inflammation (8). Our results align with this, showing a significant increase in macrophages expressing the CD14 receptor and the co-stimulatory CD40 molecule in the lungs of mice exposed to cigarette smoke for 8 weeks.

Cigarette smoke significantly impacts professional antigen-presenting cells in the lungs, such as DCs and macrophages, by promoting a pro-inflammatory state marked by the elevated production of cytokines like TNF- α , IL-12, and IL-23 (8, 11, 19). These cytokines contribute to the sustained inflammatory environment in the lungs by driving Th1 and Th17 cell-mediated lung damage (21, 22). In our study, we found that continuous exposure to cigarette smoke substantially increased the number of macrophages producing TNF- α , IL-12, and IL-23, as well as DCs expressing NLRP3 and producing IL-12 and IL-23, thereby likely contributing to the development and progression of Th1 and Th17 cell-induced inflammation in the lungs. Notably, the increased presence of IL-12 and IL-23-producing DCs was associated with a higher count of TNF- α -producing CD4⁺ and CD8⁺ T cells in the lungs of mice exposed to cigarette smoke for 8 weeks. These T cells, in a TNF- α -dependent manner, trigger apoptosis in alveolar epithelial cells, resulting in alveolar damage and compromised gas exchange (23, 24). Moreover, TNF- α produced by T cells increases the expression of E and P selectins on lung endothelial cells, facilitating the recruitment of circulating leukocytes into the lungs, which further intensifies the inflammatory response and worsens lung injury (24).

Conclusion

In conclusion, extended exposure to cigarette smoke promoted the development of an inflammatory phenotype in innate immune cells within the lungs, contributing to the progression and worsening of immune cell-driven inflammatory lung diseases.

References

1. Hikiş P, Jaceník D. The Tobacco Smoke Component, Acrolein, as a Major Culprit in Lung Diseases and Respiratory Cancers: Molecular Mechanisms of Acrolein Cytotoxic Activity. *Cells* 2023;12(6):879. [[CrossRef](#)][[PubMed](#)]
2. Szoka P, Lachowicz J, Cwiklińska M, Lukaszewicz A, Rybak A, Baranowska U, et al. Cigarette Smoke-Induced Oxidative Stress and Autophagy in Human Alveolar Epithelial Cell Line (A549 Cells). *Adv Exp Med Biol* 2019;1176:63-9. [[CrossRef](#)][[PubMed](#)]
3. Upadhyay P, Wu CW, Pham A, Zeki AA, Royer CM, Kodavanti UP, et al. Animal models and mechanisms of tobacco smoke-induced chronic obstructive pulmonary disease (COPD). *J Toxicol Environ Health B Crit Rev* 2023;26(5):275-305. [[CrossRef](#)][[PubMed](#)]
4. Casal A, Suárez-Antelo J, Riveiro V, Ferreira L, Rodríguez-Núñez N, Toubes ME, et al. Smoking-related interstitial lung disease: A narrative review. *Chron Respir Dis* 2024;21:14799731241291538. [[CrossRef](#)][[PubMed](#)]
5. Li Y, Hecht SS. Carcinogenic components of tobacco and tobacco smoke: A 2022 update. *Food Chem Toxicol* 2022;165:113179. [[CrossRef](#)][[PubMed](#)]
6. Furnari S, Emma R, Caruso M, Furneri PM, Fuochi V. Evaluating the Risks of Heated Tobacco Products: Toxicological Effects on Two Selected Respiratory Bacteria and Human Lung Cells. *Toxics* 2025;13(2):70. [[CrossRef](#)][[PubMed](#)]
7. Fan J, Zhou Y, Meng R, Tang J, Zhu J, Aldrich MC, et al. Cross-talks between gut microbiota and tobacco smoking: a two-sample Mendelian randomization study. *BMC Med* 2023;21(1):163. [[CrossRef](#)][[PubMed](#)]
8. Hikichi M, Mizumura K, Maruoka S, Gon Y. Pathogenesis of chronic obstructive pulmonary disease (COPD) induced by cigarette smoke. *J Thorac Dis* 2019;11(Suppl 17):S2129-S2140. [[CrossRef](#)][[PubMed](#)]
9. Quan DH, Kwong AJ, Hansbro PM, Britton WJ. No smoke without fire: the impact of cigarette smoking on the immune control of tuberculosis. *Eur Respir Rev* 2022;31(164):210252. [[CrossRef](#)][[PubMed](#)]
10. Caruso M, Emma R, Distefano A, Rust S, Poulas K, Giordano A, et al. Comparative assessment of electronic nicotine delivery systems aerosol and cigarette smoke on endothelial cell migration: The Replica Project. *Drug Test Anal* 2023;15(10):1164-74. [[CrossRef](#)][[PubMed](#)]
11. Kastratovic N, Markovic V, Harrell CR, Arsenijevic A, Stojanovic MD, Djonov V, et al. Effects of Combustible Cigarettes and Electronic Nicotine Delivery Systems on the Development and Progression of Chronic Lung Inflammation in Mice. *Nicotine Tob Res* 2024;26(6):704-14. [[CrossRef](#)][[PubMed](#)]
12. Kastratovic N, Zdravkovic N, Cekerevac I, Sekerus V, Harrell CR, Mladenovic V, et al. Effects of Combustible Cigarettes and Heated Tobacco Products on Systemic Inflammatory Response in Patients with Chronic Inflammatory Diseases. *Diseases* 2024;12(7):144. [[CrossRef](#)][[PubMed](#)]
13. Kastratovic N, Cekerevac I, Sekerus V, Markovic V, Arsenijevic A, Volarevic A, et al. Effects of combustible cigarettes and heated tobacco products on immune cell-driven inflammation in chronic obstructive respiratory diseases. *Toxicol Sci* 2024;200(2):265-76. [[CrossRef](#)][[PubMed](#)]
14. Strzelak A, Ratajczak A, Adamiec A, Feleszko W. Tobacco Smoke Induces and Alters Immune Responses in the Lung Triggering Inflammation, Allergy, Asthma and Other Lung Diseases: A Mechanistic Review. *Int J Environ Res Public Health* 2018;15(5):1033. [[CrossRef](#)][[PubMed](#)]
15. McGrath JJC, Vanderstocken G, Dvorkin-Gheva A, Cass SP, Afkhami S, Fantauzzi MF, et al. Cigarette smoke augments CSF3 expression in neutrophils to compromise alveolar-capillary barrier function during influenza infection. *Eur Respir J* 2022;60(2):2102049. [[CrossRef](#)][[PubMed](#)]
16. Riaz B, Sohn S. Neutrophils in Inflammatory Diseases: Unraveling the Impact of Their Derived Molecules and Heterogeneity. *Cells* 2023;12(22):2621. [[CrossRef](#)][[PubMed](#)]
17. Bergantini L, Cameli P, d'Alessandro M, Vagaggini C, Refini RM, Landi C, et al. NK and NKT-like cells in granulomatous and fibrotic lung diseases. *Clin Exp Med* 2019;19(4):487-94. [[CrossRef](#)][[PubMed](#)]
18. Malainou C, Abdin SM, Lachmann N, Matt U, Herold S. Alveolar macrophages in tissue homeostasis, inflammation, and infection: evolving concepts of therapeutic targeting. *J Clin Invest* 2023;133(19):e170501. [[CrossRef](#)][[PubMed](#)]
19. Lugg ST, Scott A, Parekh D, Naidu B, Thickett DR. Cigarette smoke exposure and alveolar macrophages: mechanisms for lung disease. *Thorax* 2022;77(1):94-101. [[CrossRef](#)][[PubMed](#)]
20. Cristaldi M, Buscetta M, Cimino M, La Mensa A, Giuffrè MR, Fiore L, et al. Caspase-8 activation by cigarette smoke induces pro-inflammatory cell death of human macrophages exposed to lipopolysaccharide. *Cell Death Dis* 2023;14(11):773. [[CrossRef](#)][[PubMed](#)]
21. Riches DWH, Martin TR. Overview of Innate Lung Immunity and Inflammation. *Methods Mol Biol* 2018;1809:17-30. [[CrossRef](#)][[PubMed](#)]
22. Misra DP, Agarwal V. Th17.1 lymphocytes: emerging players in the orchestra of immune-mediated inflammatory diseases. *Clin Rheumatol* 2022;41(8):2297-308. [[CrossRef](#)][[PubMed](#)]
23. Williams M, Todd I, Fairclough LC. The role of CD8+T lymphocytes in chronic obstructive pulmonary disease: a systematic review. *Inflamm Res* 2021;70(1):11-8. [[CrossRef](#)][[PubMed](#)]
24. Kheradmand F, Zhang Y, Corry DB. Contribution of adaptive immunity to human COPD and experimental models of emphysema. *Physiol Rev* 2023;103(2):1059-93. [[CrossRef](#)][[PubMed](#)]

Originalni rad

UDC: 612.2.017:612.014.46
doi: 10.5633/amm.2025.0418

UTICAJ PRODUŽENOG IZLAGANJA DUVANSKOM DIMU NA FENOTIP I FUNKCIJU ČELIJA UROĐENE IMUNOSTI U PLUĆIMA

Vladislav Volarević¹, Nikolina Kastratović¹, Ana Volarević²

¹Univerzitet u Kragujevcu, Fakultet medicinskih nauka, Katedra Genetika, mikrobiologija i imunologija, Centar za smanjenje štetnih efekata bioloških i hemijskih hazarda, Kragujevac, Srbija

²Univerzitet u Kragujevcu, Fakultet medicinskih nauka, Katedra Psihologija, Centar za smanjenje štetnih efekata bioloških i hemijskih hazarda, Kragujevac, Srbija

Kontakt: Vladislav Volarević
Svetožara Markovića 69, 34000 Kragujevac, Srbija
E-mail: vladislav.volarevic@medfak.ni.ac.rs

Duvanski dim sadrži različite štetne supstance koje mogu menjati fenotip i funkciju imunskih ćelija u plućima. Da bismo bolje razumeli kako produženo izlaganje duvanskom dimu utiče na ćelije urođene imunosti (neutrofile, makrofage, dendritske ćelije, prirodne ćelije ubice (engl. *natural killer* – NK) i prirodne ćelije ubice T (engl. *natural killer T* – NKT) u plućima, obavili smo istraživanje na BALB/c miševima, koji su bili izloženi duvanskom dimu četiri ili osam nedelja. Duvanski dim je generisan pomoću *LM1 Borgwaldt* mašine. Miševi iz kontrolne grupe bili su izloženi vazduhu. Korišćenjem protočne citometrije i intracelularnim bojenjem imunskih ćelija infiltriranih u plućima uočeno je da produženo izlaganje duvanskom dimu značajno povećava prisustvo inflamacijskih neutrofila koji eksprimiraju CD14 (engl. *cluster of differentiation 14* – CD14) receptor, NK i NKT ćelija koje proizvode faktor nekroze tumora- α (engl. *tumor necrosis factor* – TNF- α), kao i makrofaga koji eksprimiraju CD40 (engl. *cluster of differentiation 40* – CD40) receptor. Takođe, značajno veći broj dendritskih ćelija koje proizvode interleukin-12 (IL-12) i interleukin-23 (IL-23) primećen je u plućima miševa koji su bili izloženi duvanskom dimu osam nedelja nego u plućima životinja koje su bile izložene vazduhu. Pored toga, značajno veći broj makrofaga koji proizvode TNF- α , IL-12, IL-23, kao i značajno veći broj neutrofila i NK i NKT ćelija koje proizvode TNF- α primećen je u plućima miševa koji su bili izloženi duvanskom dimu osam nedelja nego u plućima životinja koje su bile izložene duvanskom dimu četiri nedelje. Dakle, može se zaključiti da produženo izlaganje duvanskom dimu pospešuje stvaranje inflamacijskog fenotipa u urođenim imunskim ćelijama i da tako podstiče progresiju i pogoršanje inflamacijskih oboljenja pluća.

Acta Medica Medianae 2025; 64(4): 145–152.

Ključne reči: duvanski dim, pluća, makrofagi, neutrofili, dendritske ćelije

"This work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) Licence".